




CCAMLR

Commission for the Conservation of Antarctic Marine Living Resources
Commission pour la conservation de la faune et la flore marines de l'Antarctique
Комиссия по сохранению морских живых ресурсов Антарктики
Comisión para la Conservación de los Recursos Vivos Marinos Antárticos

Fishery Report 2015: Exploratory fishery for *Dissostichus* spp. in Subarea 88.2

FISHERY REPORT



The map above shows the management areas within the CAMLR Convention Area, the specific region related to this report is shaded.

Throughout this report the CCAMLR fishing season is represented by the year in which that season ended, e.g. 2015 represents the 2014/15 CCAMLR fishing season (from 1 December 2014 to 30 November 2015).

Fishery Report 2015: Exploratory fishery for *Dissostichus* spp. in Subarea 88.2

Introduction to the fishery

1. This report describes the exploratory longline fishery for Antarctic (*Dissostichus mawsoni*) and Patagonian toothfish (*D. eleginoides*) in Subarea 88.2.
2. Statistical Subarea 88.2 is divided into nine small-scale research units (SSRUs) (A–I) (see Figure 1). Toothfish (*Dissostichus* spp.) in SSRUs 882C–H are thought to comprise a single stock and are currently managed with two catch limits: one for SSRUs 882C–G (slope/shelf SSRUs), and SSRU 882H (northern seamounts). SSRU 882I and SSRUs 882A–B have catch limits of 0 tonnes. *Dissostichus* spp. in SSRUs 882A–B are considered to be part of the Ross Sea region stock and are managed within the assessment for Subarea 88.1.
3. The limits on the exploratory fishery for *Dissostichus* spp. in Subarea 88.2 (SSRUs 882C–H) are described in Conservation Measure (CM) 41-10, the catch limits for by-catch species are defined in CMs 33-03 and 41-10.
4. In 2011, the Commission revised the boundaries of the SSRUs in Subarea 88.2 such that 76% of the yield was assigned to the region between 70°50'S and 65°00'S (redefined as SSRU 882H) and the remaining 24% of the yield was assigned to the region south of 70°50'S (SSRUs 882C–G) as outlined in SC-CAMLR-XXX, Annex 7, paragraph 6.127. SSRU 882I remained the same and has been closed to fishing.
5. In 2013 the Scientific Committee recognised that an increase in the rate of between-season tag recaptures from SSRU 882H, combined with a high incidence of within-season recaptures, suggested localised depletion in that SSRU. Also fishing in SSRUs 882C–G had been sporadic, was variable in location, and only two tagged fish had been recaptured. Because the abundance index for the stock assessment was not incorporating tag recaptures from the southern area, the Scientific Committee requested the Working Group on Statistics, Assessments and Modelling (WG-SAM) consider how an assessment of stock abundance could be developed for SSRUs 882C–G. The Scientific Committee was unable to provide consensus on advice for the catch limits in this subarea.
6. In 2014, the catch limit for SSRUs 882C–H was set at 390 tonnes and was split into 266 tonnes in SSRU 882H and 124 tonnes in SSRUs 882C–G (CCAMLR-XXXII, paragraphs 5.38 to 5.44).
7. In 2014, the Scientific Committee recalled the requests for proposals on how an assessment of stock abundance can be developed for the southern area and agreed to a two-year research plan in SSRUs 882C–H in which the catch limit for SSRU 882H was 200 tonnes, the fishing in SSRUs 882C–G was restricted to four research blocks (Figure 1) and the combined catch limit for the research blocks was 419 tonnes with no more than 200 tonnes to be taken from any one of the research blocks. Toothfish were to be tagged at the rate of 3 fish per tonne in SSRUs 882C–G and 1 fish per tonne in SSRU 882H. Recognising the different sizes of fish in the two areas, the tag-overlap statistic was to be calculated separately for each of SSRU 882H and SSRUs 882C–G (SC-CAMLR-XXXIII, paragraph 3.173).

8. Also in 2014, a plan for research to be conducted in the northern region of SSRUs 882A–B was agreed and included in CM 41-10, Annex 41-10/B. Because this research was conducted in the part of Subarea 88.2 that is included in the ‘Ross Sea region’ stock assessment, the 200 tonnes research catch allocation was set aside from the catch limit established in CM 41-09 (see Appendix 1 for details of this research plan).




Figure 1: Research blocks in Subarea 88.2 in 2015. Research blocks 882_1 to 882_4 in SSRUs D–G are as set out in CM 41-10, Annex 41-10/A, the research blocks in SSRUs 882A–B are described in CM 41-10, Annex 41-10/B.

9. In 2015, 13 vessels (from seven Members) fished in SSRUs 882C–H. For 2016, eight Members with a total of 19 vessels have notified their intention to participate in the exploratory fishery for *Dissostichus* spp. in Subarea 88.2.

10. The timing and duration of fishing activity in Subarea 88.2 has been highly variable over time. Vessels tend to begin fishing in SSRU 882H, where most of the fishing has occurred, before moving south to SSRUs 882C–G. The fishing season in Subarea 88.2 tends to peak and close a little later than in Subarea 88.1, reflecting the movement of vessels from Subarea 88.1 to Subarea 88.2 after the end of the Subarea 88.1 fishery.

Reported catch

11. The historical catches of *Dissostichus* spp. from Subarea 88.2 are provided in Table 1. In 2015, the total reported commercial catch of *Dissostichus* spp. in Subarea 88.2 (SSRUs D–H)

was 624 tonnes. This was divided between research block 882_2 (188 tonnes), 882_3 (146 tonnes), 882_4 (82 tonnes) and SSRU H (208 tonnes).

Table 1: Catch history for *Dissostichus* spp. in Subarea 88.2. (Source: STATLANT data for past seasons, and catch and effort reports for the current season, past reports for IUU catch.). Note that the STATLANT data includes catch occurring in SSRUs 882A–B as part of Subarea 88.2 catch although the stock assessment includes catch in these SSRUs as part of the Ross Sea Region stock assessment.

Season	Subarea 88.2				
	Catch limit (tonnes)	Reported catch (tonnes)			Estimated IUU catch (tonnes)
		<i>D. mawsoni</i>	<i>D. eleginoides</i>	Total	
1997	1980	0	0	0	-
1998	63	0	0	0	-
1999	0	0	0	0	-
2000	250	0	0	0	-
2001	250	0	0	0	-
2002	250	41	0	41	0
2003	375	106	0	106	0
2004	375	374	0	375	0
2005	375	411	0	411	0
2006	487	514	0	514	15
2007	547 ¹	347	0	347	0
2008	567	416	0	416	0
2009	567	484	0	484	0
2010	575	314	0	314	0
2011	575 ¹	590	0	590	*
2012	530 ¹	424	0	425	*
2013	530	475 ^q	0	476	*
2014	390	426 ^q	0	426	*
2015	819 ¹	622 ^q	0	624	*

¹ In Subarea 88.2, the catch limit includes allocation of catch for research fishing of 20 tonnes in 2007, 10 tonnes in 2011 and 2012 and 200 tonnes in 2014.

* IUU catch levels not estimated; no evidence of IUU presence or activity reported.

^q Some catch data in this year is quarantined. The following catch is not included in the reported catch table above:

2013 – vessel *Yantar 35*, 1 tonne of *D. mawsoni*

2014 – vessel *Yantar 35*, <1 tonne of *D. mawsoni*

2015 – vessel *Yantar 35*, 2 tonnes of *D. mawsoni*.

Illegal, unreported and unregulated (IUU) fishing

12. Illegal, unreported and unregulated (IUU) catch in Subarea 88.2 was estimated at 15 tonnes in 2006 taken from the south of SSRU 882A (Table 1). Following the recognition of methodological issues regarding the estimation of IUU catch levels since 2011, evidence of IUU presence or activity has continued to be recorded but no corresponding estimates of the IUU catch for *Dissostichus* spp. have been provided (SC-CAMLR-XXIX, paragraph 6.5). No evidence of IUU presence or activity has been reported in Subarea 88.2 since 2006.

Data collection

13. SSRUs 882C–H are managed under the umbrella of the exploratory fisheries CM 41-01 have an associated data collection plan (Annex 41-01/A), a research plan (Annex 41-01/B) and a tagging program (Annex 41-01/C). The data collected under this conservation measure are described below.

Biological data

14. The collection of biological data under CM 23-05 is conducted as part of the CCAMLR Scheme of International Scientific Observation. In exploratory longline fisheries targeting *D. mawsoni* and *D. eleginoides*, biological data collection includes representative samples of length, weight, sex and maturity stage, as well as collection of otoliths for age determination of the target and most frequently taken by-catch species and is most recently described in WG-FSA-15/40.

Length distributions of catches

15. The length-frequency distributions of *D. mawsoni* caught in this fishery from 2006 to 2015 are presented in Figure 2. These length-frequency distributions are unweighted (i.e. they have not been adjusted for factors such as the size of the catches from which they were collected). The interannual variability exhibited in the figure may reflect differences in the fished population but is also likely to reflect changes in the gear used, the number of vessels in the fishery and the spatial and temporal distribution of fishing.

16. The length-frequency distribution of the *D. mawsoni* catch in SSRU 882H appears to be very stable with little evidence of change in length over time (Figure 2). In SSRUs 882C–G, there was a distinct mode at about 60–80 cm, meaning that in years when there was fishing in these SSRUs there was bimodality in the overall length-frequency distribution for the subarea. Note that in SSRUs 882A–B fishing occurred in the northern region while in earlier years fishing in these SSRUs occurred in the south on the continental slope/shelf.




Figure 2: Annual length-frequency distributions of *Dissostichus mawsoni* caught in Subarea 88.2 (top panel) and in each SSRU (lower panels). The number of hauls from which fish were measured (N) and the number of fish measured (n) in each year are provided.

Tagging

17. Under CM 41-01, each longline vessel fishing in exploratory fisheries for *Dissostichus* spp. has been required to tag and release *Dissostichus* spp. according to CCAMLR tagging protocol and the required tagging rate per tonne of green weight caught specified in the fishery specific conservation measure (CM). In order to ensure that there is sufficient overlap between the length distribution of those fish that are tagged by a vessel and of all the fish that are caught by that vessel, each vessel is required to achieve a minimum tag-overlap statistic of 60% (see Annex 41-01/C, footnote 3). To avoid low sample size artefacts, the requirement for a 60% tag-overlap statistic does not apply to vessels that tag at the required rate but tag less than 30 fish (Table 2).

18. Since 2002, a total of 7 018 *D. mawsoni* and six *D. eleginoides* have been tagged and 508 and four respectively have been recaptured (Table 3 provides the details for *D. mawsoni*). Prior to 2015, the requirements for minimum tagging rate (1 fish per tonne of green weight caught) and tag overlap statistic (60%) applied across the whole of Subarea 88.2. Beginning in 2015, the minimum tagging rate was revised and applied as follows:

SSRU H	1 fish per tonne
SSRUs C, D, E, F, G	3 fish per tonne.

The minimum tag overlap statistic (60%) was also applied by the SSRU groups above.

(b)

SSRU	Required tagging rate	Flag State	Vessel name	2015
A, B	3	New Zealand	<i>Janas</i>	3.3 (72, -)
		Norway	<i>Seljevaer</i>	3.1 (61, -)
		UK	<i>Argos Froyanes</i>	3.1 (85, *)
C, D, E, F, G	3	Australia	<i>Antarctic Chieftain</i>	3.2 (85, -)
		Korea, Republic of	<i>Kostar</i>	3.2 (*, -)
		Korea, Republic of	<i>Sunstar</i>	3.2 (77, -)
		Norway	<i>Seljevaer</i>	3.1 (74, -)
		Russia	<i>Yantar 31</i>	3.0 (*, -)
		Ukraine	<i>Simeiz</i>	3.1 (83, -)
H	1	Australia	<i>Antarctic Chieftain</i>	1.1 (84, -)
		Norway	<i>Seljevaer</i>	1.0 (60, -)
		Ukraine	<i>Simeiz</i>	1.0 (69, -)

Table 3: The number of individuals of *Dissostichus mawsoni* tagged in Subarea 88.2 (a) from 2005 to 2014, and (b) since 2015. The number of fish recaptured by each vessel is provided in brackets.

(a)

Flag State	Vessel name	Season									
		2005	2006	2007	2008	2009	2010	2011	2012	2013	2014
Argentina	<i>Antartic II</i>		16 (4)	2 (1)							
	<i>Argenova XXI</i>						8 (0)				
Chile	<i>Isla Eden</i>					5 (0)					
Korea,	<i>Hong Jin No. 701</i>									7 (0)	20 (0)
Republic of	<i>Hong Jin No. 707</i>					17 (3)		40 (3)	38 (1)		22 (1)
	<i>Jung Woo No. 3</i>						6 (0)	35 (0)			
	<i>Kostar</i>									11 (0)	10 (0)
	<i>Sunstar</i>									8 (1)	33 (1)
New Zealand	<i>Antarctic Chieftain</i>					78 (0)		46 (1)	59 (9)	321 (42)	171 (19)
	<i>Avro Chieftain</i>	269 (16)			349 (33)						
	<i>Janas</i>		64 (3)			58 (2)		30 (3)	99 (17)	62 (0)	21 (0)
	<i>San Aspiring</i>	1 (0)						190 (17)			
Norway	<i>Froyanes</i>		194 (18)	97 (18)							
	<i>Seljevaer</i>									9 (1)	30 (0)
Russia	<i>Chio Maru No. 3</i>							90 (2)	101 (1)		
	<i>Gold Gate</i>							44 (16)			
	<i>Palmer</i>									55 (3)	24 (0)
	<i>Sparta</i>							50 (3)	36 (10)	12 (3)	27 (0)
	<i>Volna</i>	14 (0)		55 (2)							
	<i>Yantar</i>	77 (0)	1 (0)	100 (4)							
	<i>Yantar 31</i>									2 (0)	13 (0)
South Africa	<i>Ross Mar</i>						120 (27)				
Spain	<i>Tronio</i>						15 (2)	52 (4)			
UK	<i>Argos Froyanes</i>				38 (0)	51 (0)	250 (38)	68 (2)	210 (49)	15 (4)	67 (3)
	<i>Argos Georgia</i>		76 (0)	0 (1)		182 (21)	9 (1)	58 (13)			13 (5)
	<i>Argos Helena</i>		91 (3)	14 (7)		24 (0)					
Ukraine	<i>Simeiz</i>									4 (0)	12 (0)
Uruguay	<i>Ross Star</i>				2 (3)	53 (0)		16 (0)			
	<i>Viking Sur</i>			10 (0)							
Total		361 (16)	442 (28)	278 (33)	389 (36)	603 (55)	325 (43)	667 (60)	543 (87)	508 (54)	463 (29)

(b)

SSRU	Flag State	Vessel name	2015
A, B	New Zealand	<i>Janas</i>	165 (0)
	Norway	<i>Seljevaer</i>	33 (0)
	UK	<i>Argos Froyanes</i>	150 (0)
C, D, E, F, G	Australia	<i>Antarctic Chieftain</i>	240 (1)
	Korea, Republic of	<i>Kostar</i>	5 (0)
	Korea, Republic of	<i>Sunstar</i>	76 (0)
	Norway	<i>Seljevaer</i>	438 (19)
	Russia	<i>Yantar 31</i>	18 (0)
	Ukraine	<i>Simeiz</i>	351 (2)
H	Australia	<i>Antarctic Chieftain</i>	145 (25)
	Norway	<i>Seljevaer</i>	11 (1)
	Ukraine	<i>Simeiz</i>	64 (2)
Total			1865 (51)

Life-history parameters

Stock structure

19. The current working hypothesis regarding spawning dynamics and early life history of *D. mawsoni* in Subarea 88.2 is described in WG-SAM-14/26. Under this hypothesis, spawning takes place in the northern seamounts of SSRU 882H. As in other regions, spawning appears to take place during winter, and may extend over a period of several months. WG-FSA-12/48 showed via oceanographic drift simulations that eggs and larvae released on seamounts in SSRU 882H would be advected slowly to the east and then recruit to the slope in the eastern SSRUs 882F–G and in the western region of Subarea 88.3. As the juveniles grow in size, they move west along the shelf and slope. The fish gradually move deeper as they grow, feeding in the slope region in depths of 1 000–1 500 m, where they gain condition before moving north onto the seamounts in SSRU 882H to start the cycle again. Spawning fish appear to remain in the northern area for only one or two years based on tagging data (WG-SAM-14/27). As fishing effort in the southern region has been sporadic, no tagged fish released in SSRU 882H has been recaptured in any other area to date.

20. Analysis of the genetic diversity for *D. mawsoni* from Subareas 48.1 and 88.1 and Division 58.4.2 found weak genetic variation between the three areas (Smith and Gaffney, 2005). This differentiation is supported by oceanic gyres, which may act as juvenile retention systems, and by limited movement of tagged fish. Kuhn and Gaffney (2008) expanded the work of Smith and Gaffney (2005) by examining nuclear and mitochondrial single nucleotide polymorphisms on tissue samples collected from Subareas 48.1, 88.1 and 88.2 and Division 58.4.1. They found broadly similar results to those of the earlier studies, with some evidence for significant genetic differentiation between the three ocean sectors but limited evidence for differentiation within ocean sectors. A lack of genetic differentiation between stocks in different ocean sectors was reported by Mugue et al. (2014).

Parameter estimates

Standardised CPUE

21. Standardised catch-per-unit-effort (CPUE) analyses of *D. mawsoni* were updated for 2015 in WG-FSA-15/36. In SSRU 882H, standardised CPUE indices declined from 2003 to around 2011, then increased slightly to 2014 followed by a sharp increase in 2015 to the highest level since 2003. In SSRUs 882C–G, standardised CPUE indices show a strong increase from 2007 to 2013, before decreasing slightly in 2014 and 2015, though the uncertainty in the trend is high.

Catch at age

22. For the purposes of estimating fishery selectivity for the SSRUs 882C–H fishery, three strata are defined using *D. mawsoni* length- and age-frequency data: SSRU 882H, 882G and 882C–F (WG-FSA-14/56 and 14/57).

23. The numbers of otoliths collected by New Zealand vessels and subsequently aged are insufficient to represent the age frequency of the catch in each of the strata in every year. Where available, the otolith ages were used to construct annual area-specific age-length keys (ALKs), which were applied to the scaled length-frequency distributions for those years to produce annual catch-at-age distributions (WG-FSA-14/52).

Tag-recapture data

24. The tagging program in SSRUs 882C–H has resulted in 6 146 tagged fish released and 690 recaptured fish (WG-FSA-15/37). The higher tagging rate implemented in SSRUs 882C–G along with spatially structured fishing and higher catch limit in 2015 within the four research blocks resulted in the release of more than 1 200 tagged fish, and two recaptures from previous years from fish originally released on the slope/shelf in SSRUs 882C–H.

Parameter values

25. Estimates of natural mortality, length–mass, growth and maturity parameters for *D. mawsoni* in SSRUs 882C–H are as used in the Ross Sea assessment.

Stock assessment status

26. Two-area population models for *D. mawsoni* in the Amundsen Sea region have been developed for SSRUs 882C–H as current single-area models did not fully explain the patterns in the observed data on tag recaptures and age composition (WG-SAM-15/49). Although the hypothesised stock structure spans SSRUs 882C–H, these models were limited to data collected in SSRU 882H as there were few data available to inform estimation of biomass in SSRUs 882C–G. Additional data resulting from a two-year research plan initiated in 2015 are expected to better inform the assessment of the entire stock, including SSRUs 882C–G, in the future. Results showed that a two-area model with sex- and age-specific migrations from SSRUs 882C–G to SSRU 882H and back provided the best fits to the age and tag data collected in SSRU 882H. Furthermore, a resident population in SSRU 882H was not required to explain the patterns observed in the data, nor was annually varying or density-dependent migration.

27. Current catch limits are based on analysis of mark-recapture data undertaken in 2014 and will be reviewed in 2016.

By-catch of fish and invertebrates

Fish by-catch

28. Catch limits for by-catch species groups (macrourids, rajids and other species) are defined in CM 33-03 and provided in Table 4. Within these catch limits, the total catch of by-catch species in any SSRU or combination of SSRUs, as defined in relevant conservation measures, shall not exceed the following limits:

- skates and rays (rajids) – 5% of the catch limit of *Dissostichus* spp. or 50 tonnes, whichever is greater
- all other species combined – 20 tonnes.

29. If the by-catch of any one species is equal to, or greater than, 1 tonne in any one haul or set, then the fishing vessel must move at least 5 n miles away for a period of at least five days.

30. If the catch of *Macrourus* spp. taken by a single vessel in any two 10-day periods in a single SSRU exceeds 1 500 kg in a 10-day period and exceeds 16% of the catch of *Dissostichus* spp. in that period, the vessel shall cease fishing in that SSRU for the remainder of the season.

31. Skates thought to have a reasonable chance of survival are released at the surface in accordance with CM 33-03. The current by-catch limits and move-on rules for rajids are given in CM 33-03.

Table 4: Catch history for by-catch species (macrourids, rajids and other species), catch limits and number of rajids released alive in Subarea 88.2 (SSRUs 882C–H). Catch limits are for the whole fishery (see CM 33-03 for details). (Source: fine-scale data.)

Season	Macrourids		Rajids			Other species	
	Catch limit (tonnes)	Reported catch (tonnes)	Catch limit (tonnes)	Reported catch landed dead (tonnes)	Number released	Catch limit (tonnes)	Reported catch (tonnes)
2002	40	4	-	0	-	20	0
2003	60	18	-	0	-	140	8
2004	60	37	50	0	107	140	8
2005	60	21	50	0	-	140	3
2006	78	92	50	0	923	100	12
2007	88	54	50	0	-	100	13
2008	88	17	50	0	-	100	4
2009	90	58	50	0	265	100	14
2010	92	49	50	0	-	100	15
2011	92	52	50	0	169	100	13
2012	84	29	50	0	-	120	11
2013	84	25	50	0	-	120	8
2014	62	7	50	0	28	120	3
2015	99	19	50	1	131	120	7

Invertebrate by-catch including VME taxa

32. All Members are required to submit, within their general new (CM 21-01) and exploratory (CM 21-02) fisheries notifications, information on the known and anticipated impacts of their gear on vulnerable marine ecosystems (VMEs), including benthos and benthic communities such as seamounts, hydrothermal vents and cold-water corals. All of the VMEs in CCAMLR's VME Register are currently afforded protection through specific area closures. There have been 16 VME Risk Areas identified in SSRUs 882C–H. The locations and other details can be found at www.ccamlr.org/node/85695.

Incidental mortality of seabirds and marine mammals

Incidental mortality

33. The risk levels of birds in the fishery in Subarea 88.2 is category 1 (low) south of 65°S, category 3 (average) north of 65°S and overall is category 3 (SC-CAMLR-XXX, Annex 8, paragraph 8.1). There have been no reports of incidental mammal or bird mortalities in Subarea 88.2.

Mitigation measures

34. CM 25-02 applies to these areas and in recent years has been linked to an exemption for night setting in CM 24-02 and subject to a bird by-catch limit. Offal and other discharges are regulated under annual conservation measures (e.g. CMs 41-09 and 41-10).

Ecosystem implications and effects

35. Developments in evaluating ecosystem effects of the *D. mawsoni* fishery were discussed at the FEMA and FEMA2 Workshops (SC-CAMLR-XXVI/BG/06, paragraphs 45 to 48 and SC-CAMLR-XXVIII, Annex 4); much of the emphasis of those discussions was on the Ross Sea fishery and a summary of the outcomes can be found in the Fishery Report for Subarea 88.1.

Current management advice and conservation measures

36. The limits on the exploratory fishery for *Dissostichus* spp. in Subarea 88.2 are defined in CM 41-10. The limits in force for the forthcoming season are summarised in Table 5.

Table 5: Limits on the exploratory fishery for *Dissostichus* spp. in Subarea 88.2 in force (CM 41-10).

Element	Limit in force
Access (gear)	Limited to notified vessels using longlines
Catch limit	Precautionary catch limit for <i>Dissostichus</i> spp. is 619 tonnes for Subarea 88.2, applied as follows: SSRUs A, B and I – 0 tonnes SSRUs C, D, E, F and G – 419 tonnes total only in research blocks as defined in Annex 41-10/A with no more than 200 tonnes in any one research block SSRU H – 200 tonnes
Season	1 December to 31 August
Fishing operations	In accordance with CM 41-01 and the setting of research hauls is not required (Annex 41-01/B, paragraphs 3 and 4)
By-catch	Regulated by CMs 33-03 and 41-10
Mitigation	In accordance with CM 25-02, except paragraph 4 if requirements of CM 24-02 are met Daylight setting allowed under CM 24-02
Observers	Each vessel to carry at least two scientific observers, one of whom shall be appointed in accordance with the CCAMLR Scheme of International Scientific Observation
VMS	To be operational in accordance with CM 10-04
CDS	In accordance with CM 10-05
Research	Undertake research plan and tagging program as set out in Annexes 41-01/B and 41-01/C Toothfish tagged at a rate of at least 1 fish per tonne green weight caught.
Data	Daily and five-day catch and effort reporting under CMs 23-01 and 23-07 Haul-by-haul catch and effort data under CM 23-04 Biological data reported by the CCAMLR scientific observer
Target species	For the purposes of CMs 23-01 and 23-04, the target species is <i>Dissostichus</i> spp. and the by-catch is any species other than <i>Dissostichus</i> spp.
Environmental protection	Regulated by CMs 22-06, 22-07, 22-08 and 26-01

References

- Kuhn, K.L. and P.M. Gaffney. 2008. Population subdivision in the Antarctic toothfish (*Dissostichus mawsoni*) revealed by mitochondrial and nuclear single nucleotide polymorphisms (SNPs). *Ant. Sci.*, 20: 327–338.
- Mugue, N.S., A.F. Petrov, D.A. Zelenina, I.I. Gordeev and A.A. Sergeev. 2014. Low genetic diversity and temporal stability in the Antarctic toothfish (*Dissostichus mawsoni*) from near-continental seas of Antarctica. *CCAMLR Science*, 21: 1–9.
- Smith, P.J. and P.M. Gaffney. 2005. Low genetic diversity in the Antarctic toothfish (*Dissostichus mawsoni*) observed with mitochondrial and intron DNA markers. *CCAMLR Science*, 12: 43–51.

Research plan summary SSRUs 882 A–B north

Background

A1. New Zealand, Norway, Russia and the UK proposed to undertake research using standardised longline gear to sample the toothfish populations in the northern areas (61°S–66°S) of small-scale research units (SSRUs) 882A–B. The purpose of the research, as requested by the Scientific Committee (SC-CAMLR-XXXII, paragraph 3.76), is to characterise the local toothfish populations found there to better understand stock structure, movement patterns and improve estimation of population characteristics by Ross Sea spatial population models. Additional outcomes of the research relate to mapping the bathymetry of the fishable area, documenting relative abundance of Patagonian toothfish (*Dissostichus eleginoides*) and Antarctic toothfish (*D. mawsoni*), tagging toothfish for biomass estimation and for stock linkage studies, and collecting information on distribution, relative abundance and life history of by-catch species.

Specific Objectives

- A2.1 To utilise the expertise and experience of crew aboard Member vessels to explore and locate fishable habitat and sample toothfish in the northern region of SSRUs 882A–B (Figure A1).
- A2.2 To collect information on bathymetry in the northern region of SSRUs 882A–B to characterise the spatial extent of fishable habitat.
- A2.3 To document the spatial distribution of toothfish species in the northern area of SSRUs 882A–B, thus providing catch and biological observations to test and improve the functionality of spatial population models (SPMs) of the Ross Sea region.
- A2.4 To tag toothfish and collect biological samples to further understand toothfish movement, migration, spawning and stock linkages within Area 88.

Advice by the Scientific Committee

A3. The Scientific Committee recommended the bathymetry mapping and survey go ahead as an effort-limited ‘prospecting’ phase research design with a maximum of 6 900 hooks per set and 17 250 hooks per cluster of stations, a minimum cluster separation of 10 n miles and a total effort limit of 244 950 hooks set per vessel and a tagging rate of 3 fish per tonne of catch (SC-CAMLR-XXXIII, paragraph 3.221). The Scientific Committee agreed that an upper catch limit of 50 tonnes per vessel deducted from the catch limit from the Ross Sea region was appropriate for the scope of the research and recommended that the Commission consider

appropriate options to account for the survey catches, noting that a proposal for this purpose was submitted by New Zealand (SC-CAMLR-XXXIII/09).

A4. The results of the first year of research were presented in WG-SAM-15/31, 15/42, 15/46 and WG-FSA-15/03 and reported variable, but generally high, catch rates almost exclusively of *D. mawsoni*, with low levels of by-catch. Most fish were mature, with an age structure in each research block comparable to estimates from the Ross Sea region SPM. The vessels that participated in the survey in 2015 are shown in Table A1.

A5. Despite agreement in the Scientific Committee that the research should continue (SC-CAMLR-XXXIV, paragraphs 3.198 to 3.201), the Commission did not approve the continuation of research in 2016 (CCAMLR-XXXIV, paragraphs 5.36 to 5.44).




Figure A1: The six largest features in SSRU 882A and five largest areas in SSRU 882B with depth shallower than 2 500 m based on GEBCO bathymetry, and an 18 520 m (10 n miles) grid of points to show spatial scale. Rectangular polygons indicate the research blocks, each approximately 23 500 km² (153 km × 153 km).

Table A1: Four vessels that participated in the survey in 2015.

Members:	New Zealand	United Kingdom	Norway	Russia
Vessels 2015:	<i>Janas</i>	<i>Argos Froyanes</i>	<i>Seljevaer</i>	<i>Mys Marii</i>